

On computations of angular momentum and its flux in numerical relativity

Emanuel Gallo^{1*}, Luis Lehner^{2†} and Osvaldo M. Moreschi^{1‡}

¹Instituto de Física (CONICET)

FaMAF, Universidad Nacional de Córdoba

Ciudad Universitaria, (5000) Córdoba, Argentina.

²Department of Physics & Astronomy, LSU,

Baton Rouge, LA 70803, USA.

December 15, 2008

Abstract

The purpose of this note is to point out ambiguities that appear in the calculation of angular momentum and its radiated counterpart when some simple formulae are used to compute them. We illustrate, in two simple different examples, how incorrect results can be obtained with them. Additionally, we discuss the magnitude of possible errors in well known situations.

1 Introduction

It is well known that the notion of total angular momentum in general relativity is sensitive to the so called problem of supertranslation ambiguities. This topic involves subtleties and complications inherent in the infinite dimensional nature of the asymptotic symmetry group of asymptotically flat spacetimes, namely, the BMS group. On the other hand, there is a clear need to compute these quantities to extract valuable physical information in numerical constructed spacetimes to connect with observations of a variety of sources. A few proposals exist to do this either locally[1, 2, 3, 4, 5] or asymptotically[1, 6, 7, 8, 9, 10, 11, 12]¹. The dependence of any notion of relativistic angular momentum on supertranslations has been the matter of numerous studies. The problem is complex and, unfortunately, most definitions of angular momentum either show supertranslation ambiguities or require of additional structure (at the expense of reducing generality in their application).

In spite of all these difficulties, the need for tackling concrete calculations of angular momentum in numerically constructed spacetimes has motivated the use of simple and practical formulae for the computation of angular momentum. These formulae lend themselves for a straightforward computation and, in relatively simple scenarios, have given the expected answers. One option that has been employed in several works has been suggested in [13], where it is proposed to compute the angular momentum from the expression

$$J_{[i]} = \frac{1}{16\pi} \text{Re} \left\{ \oint_{\Sigma^+} dS \Phi_{[i]} \left[2\Psi_1^0 - 2\sigma^0 \bar{\sigma}^0 - \bar{\sigma}(\sigma^0 \bar{\sigma}^0) \right] \right\}; \quad (1)$$

which is a truncated version of the one defined in [8]; in fact, a supermomentum part is missing. As mentioned above, supertranslations issues can obscure angular momentum definitions (and expressions derived from them) and indeed, as we show in the next sections, eq. (1) is susceptible to them as well.

We discuss in the next examples two simple cases where expression (1) gives an incorrect result.

*Member of CONICET; email: egallo@famaf.unc.edu.ar.

†email: lehner@phys.lsu.edu

‡Member of CONICET; email: moreschi@fis.uncor.edu.

¹We are not presenting an exhaustive list of references, since this is not a review paper on angular momentum. We apologize for our arbitrary choice of references.

2 Examples

2.1 A supertranslated and boosted Schwarzschild

Let us consider, for simplicity, the case of a spacetime describing two orbiting black holes which after merger give rise to a non-spinning black hole. Furthermore, we consider a case where the final black hole has a kick. This scenario, can easily be realised with a non-equal mass binary with individual spins adjusted such that the final spin is zero². Therefore, the black hole will generically appear in the grid which is supertranslated from that where the black hole would be at rest.

To simplify the analysis, we further assume that a natural coordinate system can be constructed such that, asymptotically, the system can be surrounded by round spheres; that is, one can construct a Bondi type coordinate system from it. The delicate issue is that this inertial asymptotic coordinate system results, generically, supertranslated and boosted with respect to the intrinsic asymptotic coordinate system which has the spherical symmetry of the remaining black hole. This in turn, implies that the asymptotic fields acquire a non-trivial dependence on the supertranslation and boost. To compare what would be obtained in both systems, we denote with primes the asymptotic coordinate system with spherical symmetry and unprimed that naturally constructed in the numerical implementation; then one would have the BMS transformation

$$u = K(\zeta', \bar{\zeta}') (u' - \gamma(\zeta', \bar{\zeta}')) + \frac{u_1(u', \zeta', \bar{\zeta}')}{r'} + O\left(\frac{1}{r^2}\right), \quad (2)$$

$$r = \frac{r'}{K(\zeta', \bar{\zeta}')} + O(r^0), \quad (3)$$

$$\zeta = \frac{a\zeta' + b}{c\zeta' + d} + O\left(\frac{1}{r'}\right); \quad (4)$$

where K is the Lorentz factor associated with the boost, γ determines the supertranslation and (a, b, c, d) are complex constants which determine the boost; for details see [19].

In what follows it will be convenient to use the notation

$$e^{2i\delta} = \frac{\bar{\zeta}_{\bar{\zeta}'}}{\zeta_{\zeta'}} = \left(\frac{c\zeta' + d}{\bar{c}\bar{\zeta}' + \bar{d}} \right)^2. \quad (5)$$

Although the calculation of expression (1) on a spherically symmetric section gives the expected null result we will show next that it gives a non-zero result when calculated at a supertranslated and boosted section.

As explained elsewhere[20, 21] the above transformation implies a transformation of the null frames, and in particular, the components of the Weyl tensor and the asymptotic shear are affected. More concretely, one has

$$\Psi_1^0 = \frac{e^{i\delta}}{K^3} \left(\Psi_1'^0 - 3 \frac{\bar{\partial}' u}{K} \Psi_2'^0 + 3 \left(\frac{\bar{\partial}' u}{K} \right)^2 \Psi_3'^0 + \left(\frac{\bar{\partial}' u}{K} \right)^3 \Psi_4'^0 \right). \quad (6)$$

Only $\Psi_2'^0 = -m$ is different from zero in the spherically symmetric frame; therefore one has

$$\Psi_1^0 = \frac{3m}{K^4} e^{i\delta} \bar{\partial}' u; \quad (7)$$

where m is the mass of the resulting black hole. Similarly the shear is

$$\sigma^0 = -\frac{1}{K} e^{i2\delta} \bar{\partial}'^2 \gamma. \quad (8)$$

Let us now calculate expression (1) in detail, at a $u = \text{const.}$ section, and discuss what is obtained. First, note that the factors $\Phi_{[i]}$ in the integrand, are spin weighted quantities with spin weight -1. Therefore they can be expressed in terms of an edth bar operator acting on a spin weight 0 quantity,

²A straightforward calculation can be employed to adjust the physical parameters so that this scenario is realized as discussed by the simple model [14] or the fitted expansions presented in [15, 16]. The resulting kick velocity can be estimated by suitable fits as in [15, 17, 18].

namely $\Phi_{[i]} = \bar{\partial}a_{[i]}$; where, to pick the angular momentum components $a_{[i]}$ should be imaginary[11]. Consequently, each $a_{[i]}$ is a quantity of spin weight 0 satisfying $\bar{\partial}\bar{\partial}a_{[i]} = -a_{[i]}$.

We should now express the integration appearing in (1) in terms of the primed, spherically symmetric Bondi system. Let us observe that for all these quantities $a_{[i]}$, due to the transformation properties of the edth operator[22], one has

$$\bar{\partial}a = \frac{e^{-i\delta}}{K}\bar{\partial}'a'; \quad (9)$$

where $a' = a'(\zeta', \bar{\zeta}')$ is understood in terms of the primed angular coordinates. Next, one must also take into account that $dS = K^2 dS'$, then the first term of (1), involving Ψ_1^0 , is

$$\Re \left\{ \oint_{\Sigma^+} \Phi_{[i]} \Psi_1^0 dS \right\} = \Re \left\{ \oint_{\Sigma^+} \frac{e^{-i\delta}}{K} \Phi'_{[i]} \frac{3m}{K^4} e^{i\delta} \bar{\partial}' [K(u' - \gamma)] K^2 dS' \right\}. \quad (10)$$

In order to simplify the discussion we can assume that the calculation is performed at the section $u = 0$, which is equivalent to $u' = \gamma$; therefore, one has $\bar{\partial}' [K(u' - \gamma)] = (u' - \gamma)\bar{\partial}' K - K\bar{\partial}' \gamma = -K\bar{\partial}' \gamma$; which implies

$$\Re \left\{ \oint_{\Sigma^+} \Phi_{[i]} \Psi_1^0 dS \right\} = -\Re \left\{ 3m \oint_{\Sigma^+} \Phi'_{[i]} K^{-2} \bar{\partial}' \gamma dS' \right\}. \quad (11)$$

Since the factor $K^{-2}\bar{\partial}' \gamma$ has spin weight 1, one again can think of a spin-weight 0 potential A such that $\bar{\partial}' A = K^{-2}\bar{\partial}' \gamma$. Therefore, if the quantity A were real, this term would not have any contribution, since one would have to evaluate

$$\begin{aligned} \Re \left\{ \oint_{\Sigma^+} \Phi_{[i]} \Psi_1^0 dS \right\} &= -\Re \left\{ 3m \oint_{\Sigma^+} \bar{\partial}' a'_{[i]} \bar{\partial}' A dS' \right\} = \Re \left\{ 3m \oint_{\Sigma^+} \bar{\partial}' \bar{\partial}' a'_{[i]} A dS' \right\} \\ &= -\Re \left\{ 3m \oint_{\Sigma^+} a'_{[i]} A dS' \right\} = 0; \end{aligned} \quad (12)$$

since $a'_{[i]}$'s are imaginary and we are assuming at the moment that A is real. Unfortunately, it is not difficult to prove that actually A has an imaginary part; and furthermore, this imaginary part is only zero when the gradients of $\frac{1}{K^2}$ and γ are proportional; something that is not true in our case.

To see this, we next study the equation $\bar{\partial}A = v^2 \bar{\partial}\gamma$; where $v = K^{-1}$, and assuming A is a real quantity we will reach a contradiction. Notice that for a real spin weight zero field A , $\text{Im} [\bar{\partial}\bar{\partial}A] = 0$; since $\bar{\partial}\bar{\partial}$ is a real operator when acting on a spin weight zero quantity. This last relation can be written in terms of v and γ as

$$\bar{\partial}v^2 \bar{\partial}\gamma - \bar{\partial}\gamma \bar{\partial}v^2 = 0. \quad (13)$$

Let us denote the gradients on the sphere of the quantities v^2 and γ by $\nabla_a v^2$ and $\nabla_b \gamma$. Then the ‘cross product’ of these vectorial quantities is defined by $\epsilon^{ab} \nabla_a v^2 \nabla_b \gamma$; where ϵ^{ab} is the surface element of the unit sphere. At this point it is convenient to express the gradients in terms of the edth operator; for example for v^2 one has

$$\nabla_a v^2 = -(\bar{\partial}v^2)\bar{m}_a - (\bar{\partial}v^2)m_a; \quad (14)$$

and similarly for γ . Then the cross product is

$$\epsilon^{ab} \nabla_a v^2 \nabla_b \gamma = (\bar{\partial}v^2 \bar{\partial}\gamma - \bar{\partial}\gamma \bar{\partial}v^2) \epsilon^{ab} m_a \bar{m}_b; \quad (15)$$

which should be zero if A were real. But this is true only if the gradients are proportional. Consequently the term (11) has a non-zero contribution to the quantity $J_{[i]}$.

We now turn our attention to the second term in expression (1) which involves quadratic terms in σ . One could also in this case think of a potential X such that $\sigma^0 = \bar{\partial}'^2 X$. Then, one can prove that if X were real, then the second term of (1) involving the σ 's would not contribute to the integral[23]. However, following a similar line of arguments as above, one can prove that X will have an imaginary part if

$$-\frac{1}{K} e^{i2\delta} \neq 1, \quad (16)$$

–recall this is the factor appearing in (8)–.

Therefore, this second term will also have a non-zero contribution to the quantity $J_{[i]}$; and since it includes independent information not contained in the first term it can not cancel the first contribution.

Let us discuss further what we have found up to this point. If the adopted coordinate system, for instance the one most naturally constructed from a numerical simulation, results supertranslated and boosted with respect to the final frame adapted to the spherically symmetric final black hole an incorrect result will be obtained for the angular momentum of the system. Notice that this conclusion is more subtle than the obvious ‘Newtonian’-type observation stating that the angular momentum calculated at frames translated with respect to each other are different. In fact, even if the coordinate system is only boosted and supertranslated *—without a translation—*, with respect to the one adapted to the spherically symmetric final black hole then $J_{[i]}$ turns out to be different from zero in the Schwarzschild geometry. This is to be contrasted with what would be the case within special relativity paradigm where the absence of a translation does not affect the angular momentum calculation since the Lorentz transformation of a zero tensor is a zero tensor.

2.2 A radiating second order solution

As a second example let us consider the spacetime given by the metric

$$ds^2 = \left(-2 \frac{\dot{V}}{V} r + K_V - 2 \frac{M}{r} \right) du^2 + 2 du dr - \frac{r^2}{V^2 P_0^2} d\zeta d\bar{\zeta}; \quad (17)$$

where a dot denotes a derivative with respect to u , K_V is the Gaussian curvature, given by

$$K_V = 2V \bar{\partial} \partial V - 2\partial V \bar{\partial} V + V^2, \quad (18)$$

of the 2-metric

$$dS^2 = \frac{1}{V^2 P_0^2} d\zeta d\bar{\zeta}; \quad (19)$$

and $P_0(\zeta, \bar{\zeta})$ is the conformal factor of the unit sphere. This metric was used by Robinson and Trautman to characterize spacetimes satisfying Einstein equations admitting a congruence of null geodesic without shear and twist[24].

The scalar $V(u, \zeta, \bar{\zeta})$ can be expressed in the form $V = 1 + \Delta(u, \zeta, \bar{\zeta})$; so that $\Delta = 0$ would correspond to the Schwarzschild spacetime. The function Δ is required to satisfy the equation

$$-3 M \dot{\Delta} = (1 + 4\Delta) \bar{\partial}^2 \partial^2 \Delta - \bar{\partial}^2 \Delta \bar{\partial}^2 \Delta. \quad (20)$$

This spacetime is known as a Robinson-Trautman geometry[25]. By requiring (20) one obtains a solution of vacuum Einstein equations in second order in Δ . The Robinson-Trautman spacetimes are understood as representing a central object which has been perturbed and are known to decay in time to the Schwarzschild geometry.

Suppose one has obtained this spacetime and would like to calculate the quantity expressed in (1) at the section $u = u_0$. Then, since the asymptotic coordinate system $(u, \zeta, \bar{\zeta})$ is not inertial (not Bondi); one should look for the transformation to a Bondi frame. Let us choose a Bondi coordinate system $(\tilde{u}, \tilde{\zeta}, \tilde{\bar{\zeta}})$ such that $\tilde{u} = 0$ coincide with the section $u = u_0$. With this choice one has that on this particular section the Bondi shear is zero; $\bar{\sigma}^0 = 0$ [21]. Furthermore, since on this section the null frames are proportional, one also has $\bar{\Psi}_1^0 = 0$. Therefore in this section one obtains $J_{[i]} = 0$.

However in any other Bondi section given by $\tilde{u} \neq 0$ one will obtain $J_{[i]} \neq 0$. This can be seen due to two reasons. First one can notice that the shear with respect to the Bondi system is not stationary; in fact one can prove that $\bar{\sigma}_{,\tilde{u}} = (\bar{\partial}^2 V) V^{-1}$ [25]. One can convince oneself that $\bar{\sigma}$ can not be expressed in terms of a real potential X , that is, $\bar{\sigma} = \bar{\partial}^2 X$ as studied in the previous section. In other words such an X has an imaginary term. Therefore one deduces that there is always a flux, which in general will include the time variation of $J_{[i]}$, depending on the arbitrary initial data V . On the other hand one can also infer that $\bar{\Psi}_1^0$ will also be different from zero, at any other Bondi section; due again to the existence of radiation.

Notice that $J_{[i]} \neq 0$ at any other Bondi section $\tilde{u} \neq 0$ can also be deduced by taking (Bondi) time derivatives of $J_{[i]}$. One can see that at $\tilde{u} = 0$, one has $J_{[i],\tilde{u}} = 0$, $J_{[i],\tilde{u}\tilde{u}} = 0$ but $J_{[i],\tilde{u}\tilde{u}\tilde{u}} \neq 0$.

The weaknesses of formula (1) to calculate the angular momentum also give rise to odd results. Notice that the asymptotic value of $J_{[i]}$, for $\tilde{u} \rightarrow \infty$ is zero; since in this regime, $\tilde{u}$ will be supertranslated with respect to the $(u, \zeta, \bar{\zeta})$ frame; but not boosted (see the previous section). So, in this case one would have the curious situation that the initial value of $J_{[i]}$ is zero, a non-zero flux to the future of the initial section

and an ending with a zero value for $J_{[i]}$ in the asymptotic regime. This alone makes it difficult to identify $J_{[i]}$ with a physical notion of intrinsic angular momentum.

One instead could use the Bondi system $(u^*, \zeta^*, \bar{\zeta}^*)$ which is defined so as to coincide asymptotically with the R-T system $(u, \zeta, \bar{\zeta})$ in the regime $u \rightarrow \infty$. Then one could calculate the quantity $J_{[i]}$ for any $u^* = \text{constant}$. In this case one would obtain that in the limit $u^* \rightarrow \infty$, $J_{[i]}$ would vanish. However for any finite $u^* = \text{const.}$, the evaluation of $J_{[i]}$ would give a non zero value.

This again stresses the difficulties that one encounters in trying to use the expression (1) to obtain intrinsic information of the central object.

3 Estimate of the error in spin calculations

In order to provide a different perspective to the quantity $J_{[i]}$, in the next subsection we recall an unambiguous definition of total intrinsic angular momentum at future null infinity. Actually since this is the only definition we know of that solves simultaneously the supertranslations ambiguities for center of mass and intrinsic angular momentum[11] –without assuming further structure–; we will use it as reference for comparison.

3.1 An unambiguous definition of intrinsic angular momentum

Arguably one of the most valuable physical information aimed to be extracted from a given spacetime is that of *intrinsic angular momentum* as it relates directly with observable signatures at both gravitational and electromagnetic signals produced in a number of systems. Tied to defining such quantity is the problem of defining the center of mass of the system with respect to which the intrinsic angular momentum can be defined; consequently, the definition of both concepts must come together. A solution of these difficulties has been formulated in [11]; where the expression to calculate the charges associated to BMS generators is given by,

$$Q_{S_{cm}}(w) = \Re \left\{ 8 \int_{S_{cm}} \left(-w_2 (\Psi_1^0 + 2\sigma_0 \bar{\partial} \bar{\sigma}_0 + \bar{\partial} (\sigma_0 \bar{\sigma}_0)) + 2w_1 (\Psi_2^0 + \sigma_0 \dot{\bar{\sigma}}_0 + \bar{\partial}^2 \bar{\sigma}_0) \right) dS^2 \right\}; \quad (21)$$

where w_1 and w_2 are components of the two form w that is determined by the particular generator of BMS transformations. As with the expression appearing in [8], one can use this type of expression to calculate the total momentum, total supermomentum and total angular momentum. However, the notion of *intrinsic angular momentum* can only be obtained if one calculates the angular momentum on the *center of mass sections*, here denoted by S_{cm} .

If one calculates the angular momentum in a radiating spacetime at any other section, one would obtain a quantity with angular momentum reminiscence but with unclear intrinsic physical meaning.

3.2 How important are in practice these considerations in the computation of angular momentum?

We want to estimate now the errors in the value of angular momentum by using an unfortunate choice of frame and section. To do so, we want to have an estimate of how supertranslated are the “center of mass” sections with respect to the sections that are adapted to the coordinates used in numerical computations.

The astrophysical model that we have in mind is similar to the system that was presented in subsection 2.1, i.e, we have some compact objects (for example a binary system) that are undergoing a merge, and where the coordinate grid is in some way following the system until the moment of merger. After such event, in general the system will end up with a unique boosted (quasi)-stationary compact object (in the case of black hole collision, we will finish with something similar to a boosted Schwarzschild or Kerr black hole). Then, the new “center of mass” sections will be supertranslated and boosted with respect to the originals (those adapted to the numerical grid).

Sections that characterise rest frames are known as ‘nice’ sections[26]. The prescription to find rest frame sections uses a particular notion of supermomentum.

The supermomentum at S is defined[26] by

$$P_{lm}(S) \equiv -\frac{1}{\sqrt{4\pi}} \int_S Y_{lm}(\zeta, \bar{\zeta}) \Psi dS^2; \quad (22)$$

where $\Psi = \Psi_2 + \sigma\dot{\sigma} + \bar{\sigma}^2\bar{\sigma}$ and Y_{lm} are spherical harmonics.

Given another section $\tilde{S}$ of future null infinity, one can find another Bondi system $(\tilde{u}, \tilde{\zeta}, \tilde{\bar{\zeta}})$ such that $\tilde{S}$ is determined by $\tilde{u} = 0$. The relation between the new and the original Bondi system is given by a BMS transformation, as given by (2), (3) and (4); where now we use tildes for the new coordinates and un-tilde for the original coordinates. Note that $\tilde{S}$ can also be determined by $u = \gamma(\zeta, \bar{\zeta})$.

The section $\tilde{S}$ is said to be of type nice[26] if all the ‘spacelike’ components of the supermomentum, when calculated with respect to the adapted Bondi system, are zero;

$$\tilde{P}_{lm}(\tilde{S}) = 0 \quad \text{for } l \neq 0; \quad (23)$$

and therefore the only non-vanishing one, $\tilde{P}_{00}(\tilde{S})$, coincides with the total Bondi mass at $\tilde{S}$.

One can prove that Ψ transforms under a BMS transformation as

$$\tilde{\Psi} = \frac{1}{K^3} (\Psi - \bar{\sigma}^2 \bar{\sigma}^2 \gamma). \quad (24)$$

Then, for a section to correspond to a ‘nice’ section it must obey $\tilde{\Psi} = \text{constant}$ as otherwise some moments of P_{lm} , for $l \neq 0$, will be non-zero. The nice section equation can be understood as a condition for $\gamma(\zeta, \bar{\zeta})$ and K .

It was indicated in ref. [26] that equation (24) can be expressed by

$$\bar{\sigma}^2 \bar{\sigma}^2 \gamma = \int_0^\gamma \dot{\Psi}(u', \zeta, \bar{\zeta}) du' + \Psi(u = 0, \zeta, \bar{\zeta}) + K(\gamma; \zeta, \bar{\zeta})^3 M(\gamma). \quad (25)$$

where $M(\gamma)$ is the total mass at the section $u = \gamma$ and $K(\gamma; \zeta, \bar{\zeta})$ is the conformal factor of the BMS transformation that aligns its timelike generator (defined as $\tilde{p}_{00} = Y_{00} \frac{\partial}{\partial u}$ [26]) with the Bondi momentum at $u = \gamma(\zeta, \bar{\zeta})$.

Note that $\tilde{\Psi} = \dot{\sigma}^0 \dot{\sigma}^0$, and then it is proportional to the content of gravitational radiation. Now, let us introduce a parameter λ to measure this content of radiation, i.e., let the integrated flux, appearing in the first term of the right hand side of (25) be of order λ .

Suppose then, that we start by describing our compact bodies system with sections that are close to ‘nice’ sections. Then, after the merger, a new compact object is obtained, and due to the emission of gravitational radiation, the new ‘nice’ sections that ‘follow’ the system will, in general, not only be boosted but also supertranslated by a quantity γ . Now, let v be the kick velocity of the final compact object; then, one can observe that $K^3 \approx 1 + O(v)$. On the other hand, by computing the flux of the Bondi momentum expression we can see that

$$v \approx O\left(\frac{\lambda}{M}\right); \quad (26)$$

and $\Psi(u = 0, \zeta, \bar{\zeta}) \approx -M$, then

$$\bar{\sigma}^2 \bar{\sigma}^2 \gamma \approx O(\lambda) + O(vM) \approx O(vM). \quad (27)$$

From these estimates, and recalling that $\Psi_2^0 \approx O(M)$, and using eq.(7), we have that $\Delta\Psi_1^0 \approx O(vM^2)$.

This will be the order of the error that one would have in the computation of angular momentum by not adopting the correct sections. We conclude then, that the errors will be, at least, as large as the relativistic velocities of the astrophysical system, i.e., $\Delta J_{[i]} \approx O(vM^2)$. Consequently one can understand why the use of the flux of eq. (1), as calculated in [13], employed in several works [27, 28, 29, 30, 31] (where the velocities of the astrophysical systems under consideration are low, and therefore the error estimate $\Delta J_{[i]}$ is small) can produce sensible results. However one should be cautious with the use of formulas like (1) in cases where the individual objects move relativistically (as in the studies of high speed collisions [32]), large production of gravitational radiation or producing an object with a small final spin (where the errors described above will be important).

Finally, we stress that all these considerations are true if one assumes that the numerical grid and gauge gives rise to an extraction frame/coordinate system which is of Bondi type (which in general will not be the case). If it is not of this type, one must consider extra gauge effects like those reported in [20] and [21] in the computation of radiation flux and total momentum respectively.

4 Final comments

The difficulties shown above are related to the nature of the notion of angular momentum. In special relativity the angular momentum is expressed in terms of an antisymmetric tensor

$$J^{ab} = S^{ab} + P^a R^b - R^a P^b; \quad (28)$$

where P^a is the total momentum, S^{ab} is orthogonal to the momentum and R^a shows the dependence of the angular momentum with respect to some reference point, or origin.

The tensor S^{ab} does not depend on the choice of origin and therefore is the intrinsic angular momentum. At the center of mass line one has $J^{ab} = S^{ab}$ and the original expression of the angular momentum gives the intrinsic value.

The expression (1) does not have this translation/supertranslation dependence; and therefore its usage is ambiguous since there is no prescription of what a center of mass is. Instead, we have shown that one could use a definition of angular momentum[11] that does show this translation/supertranslation dependence; and furthermore, it can be used to define center of mass sections, in which the calculation of angular momentum gives an unambiguous intrinsic angular momentum. In particular the application of this intrinsic angular momentum to the two examples presented above, gives zero at any center of mass section.

Incidentally, note also that for general spacetimes, inferring how much angular momentum is radiated by differentiating with respect to time eq. (1) will give results strongly affected by the issues discussed here.

References

- [1] Arthur Komar. Covariant conservation laws in general relativity. *Phys. Rev.*, 113:934–936, 1959.
- [2] R. Penrose. Quasi-local mass and angular momentum in general relativity. *Proc. R. Soc. Lond. A*, 381:53–63, 1982.
- [3] László B. Szabados. Quasi-local energy-momentum and angular momentum in gr: A review article. *Living Rev. Relativity*, 7:1–135, 2004. <http://relativity.livingreviews.org/Articles/lrr-2004-4>
- [4] A. Ashtekar and B. Krishnan. Isolated and dynamical horizons and their applications. *Living Rev. Rel.*, 7:10, 2004.
- [5] S. A. Hayward. Angular momentum conservation for dynamical black holes. *Phys. Rev. D*, 74:104013, 2006.
- [6] B.D. Bramson. Relativistic angular momentum for asymptotically flat Einstein-Maxwell manifolds. *Proc. R. Soc. Lond. A*, 341:463–490, 1975.
- [7] C.R. Prior. Angular momentum in general relativity. *Proc. R. Soc. Lond.*, A(354):379, 1977.
- [8] J. Winicour. Angular momentum in general relativity. In A. Held, editor, *General Relativity and Gravitation*, volume 2, pages 71–96. Plenum, 1980.
- [9] Tevian Dray and Michael Streubel. Angular momentum at null infinity. *Class. Quantum Grav.*, 1:15–26, 1984.
- [10] Anthony Rizzi. Angular momentum in general relativity: The definition at null infinity includes the spatial definition as a special case. *Phys.Rev.D*, 63(104002), 2001.
- [11] Osvaldo M. Moreschi. Intrinsic angular momentum and center of mass in general relativity. *Class.Quantum Grav.*, 21:5409–5425, 2004.
- [12] László B. Szabados. Total angular momentum from Dirac eigenspinors. *Class.Quantum.Grav.*, 25:025007–25, 2008.
- [13] Carlos O. Lousto and Yosef Zlochower. Practical formula for the radiated angular momentum. *Phys.Rev.D.*, 76:041502(R)–4, 2007.

- [14] A. Buonanno, L. E. Kidder, and L. Lehner. Estimating the final spin of a binary black hole coalescence. *Phys.Rev.D*, 77:026004, 2008.
- [15] L. Boyle, M. Kesden and S. Nissanke, Binary black hole merger: symmetry and the spin expansion, *Phys. Rev. Lett.* **100**, 151101, 2008.
- [16] L. Rezzolla, P. Diener, E. N. Dorband, D. Pollney, C. Reisswig, E. Schnetter and J. Seiler, The final spin from the coalescence of aligned-spin black-hole binaries, *Astrophys. J.* **674**, L29, 2008.
- [17] J. G. Baker, W. D. Boggs, J. Centrella, B. J. Kelly, S. T. McWilliams, M. C. Miller and J. R. van Meter, Modeling kicks from the merger of generic black-hole binaries, *Astrophys. J.* **682**, L29, 2008.
- [18] M. Campanelli, C. O. Lousto, Y. Zlochower and D. Merritt, Large Merger Recoils and Spin Flips From Generic Black-Hole Binaries, *Astrophys. J.* **659**, L5, 2007.
- [19] Osvaldo M. Moreschi. On angular momentum at future null infinity. *Class. Quantum Grav.*, 3:503–525, 1986.
- [20] Luis Lehner and Osvaldo M. Moreschi. Dealing with delicate issues in waveforms calculations. *Phys.Rev.D*, 76:124040–1–12, 2007.
- [21] Emanuel Gallo, Luis Lehner, and Osvaldo Moreschi. Estimating total momentum at finite distances. *Phys. Rev.D*, 78:084027,2008.
- [22] E. T. Newman and R. Penrose. Note on the Bondi-Metzner-Sachs group. *J. Math. Phys.*, 7(5):863–870, 1966.
- [23] B.D. Bramson. The invariance of spin. *Proc. R. Soc. Lond. A*, 364:383–392, 1976.
- [24] I. Robinson and R. Trautman. Some spherical gravitational waves in general relativity. *Proc. R. Soc. A*, 265:463–473, 1962.
- [25] Sergio Dain, Osvaldo M. Moreschi, and Reinaldo J. Gleiser. Photon rockets and the Robinson-Trautman geometries. *Class. Quantum Grav.*, 13(5):1155–1160, 1996.
- [26] Osvaldo M. Moreschi. Supercenter of mass system at future null infinity. *Class. Quantum Grav.*, 5:423–435, 1988.
- [27] John G. Baker et al. Mergers of non-spinning black-hole binaries: Gravitational radiation characteristics. *Phys. Rev.D*, D78:044046,2008
- [28] Manuela Campanelli, Carlos O. Lousto, Hiroyuki Nakano, and Yosef Zlochower. Comparison of Numerical and Post-Newtonian Waveforms for Generic Precessing Black-Hole Binaries. arXiv:0808.0713 [gr-qc], 2008.
- [29] Carlos O. Lousto and Yosef Zlochower. Modeling gravitational recoil from precessing highly- spinning unequal-mass black-hole binaries. arXiv:0805.0159 [gr-qc], 2008.
- [30] Sergio Dain, Carlos O. Lousto, and Yosef Zlochower. Extra-Large Remnant Recoil Velocities and Spins from Near- Extremal-Bowen-York-Spin Black-Hole Binaries. *Phys. Rev.*, D78:024039, 2008.
- [31] Carlos O. Lousto and Yosef Zlochower. Further insight into gravitational recoil. *Phys. Rev.*, D77:044028, 2008.
- [32] U. Sperhake, V. Cardoso, F. Pretorius, E. Berti, and J. A. Gonzalez. The high-energy collision of two black holes. *Phys.Rev.Lett.*, 101:161101,2008